

Cortical Connectivity Moderators of Antidepressant vs Placebo Treatment Response in Major Depressive Disorder

Secondary Analysis of a Randomized Clinical Trial

Camarin E. Rolle, BS; Gregory A. Fonzo, PhD; Wei Wu, PhD; Russ Toll, PhD; Manish K. Jha, MD; Crystal Cooper, PhD; Cherise Chin-Fatt, PhD; Diego A. Pizzagalli, PhD; Joseph M. Trombello, PhD; Thilo Deckersbach, PhD; Maurizio Fava, MD; Myrna M. Weissman, PhD; Madhukar H. Trivedi, MD; Amit Etkin, MD, PhD

IMPORTANCE Despite the widespread awareness of functional magnetic resonance imaging findings suggesting a role for cortical connectivity networks in treatment selection for major depressive disorder, its clinical utility remains limited. Recent methodological advances have revealed functional magnetic resonance imaging–like connectivity networks using electroencephalography (EEG), a tool more easily implemented in clinical practice.

OBJECTIVE To determine whether EEG connectivity could reveal neural moderators of antidepressant treatment.

DESIGN, SETTING, AND PARTICIPANTS In this nonprespecified secondary analysis, data were analyzed from the Establishing Moderators and Biosignatures of Antidepressant Response in Clinic Care study, a placebo-controlled, double-blinded randomized clinical trial. Recruitment began July 29, 2011, and was completed December 15, 2015. A random sample of 221 outpatients with depression aged 18 to 65 years who were not taking medication for depression was recruited and assessed at 4 clinical sites. Analysis was performed on an intent-to-treat basis. Statistical analysis was performed from November 16, 2018, to May 23, 2019.

INTERVENTIONS Patients received either the selective serotonin reuptake inhibitor sertraline hydrochloride or placebo for 8 weeks.

MAIN OUTCOMES AND MEASURES Electroencephalographic orthogonalized power envelope connectivity analyses were applied to resting-state EEG data. Intent-to-treat prediction linear mixed models were used to determine which pretreatment connectivity patterns were associated with response to sertraline vs placebo. The primary clinical outcome was the total score on the 17-item Hamilton Rating Scale for Depression, administered at each study visit.

RESULTS Of the participants recruited, 9 withdrew after first dose owing to reported adverse effects, and 221 participants (150 women; mean [SD] age, 37.8 [12.7] years) underwent EEG recordings and had high-quality pretreatment EEG data. After correction for multiple comparisons, connectome-wide analyses revealed moderation by connections within and between widespread cortical regions—most prominently parietal—for both the antidepressant and placebo groups. Greater alpha-band and lower gamma-band connectivity predicted better placebo outcomes and worse antidepressant outcomes. Lower connectivity levels in these moderating connections were associated with higher levels of anhedonia. Connectivity features that moderate treatment response differentially by treatment group were distinct from connectivity features that change from baseline to 1 week into treatment. The group mean (SD) score on the 17-item Hamilton Rating Scale for Depression was 18.35 (4.58) at baseline and 26.14 (30.37) across all time points.

CONCLUSIONS AND RELEVANCE These findings establish the utility of EEG-based network functional connectivity analyses for differentiating between responses to an antidepressant vs placebo. A role emerged for parietal cortical regions in predicting placebo outcome. From a treatment perspective, capitalizing on the therapeutic components leading to placebo response differentially from antidepressant response should provide an alternative direction toward establishing a placebo signature in clinical trials, thereby enhancing the signal detection in randomized clinical trials.

TRIAL REGISTRATION ClinicalTrials.gov identifier: [NCT01407094](https://clinicaltrials.gov/ct2/show/study/NCT01407094)

JAMA Psychiatry. 2020;77(4):397-408. doi:10.1001/jamapsychiatry.2019.3867
Published online January 2, 2020.

← Editorial page 347

+ Supplemental content

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Amit Etkin, MD, PhD, Department of Psychiatry and Behavioral Sciences, Stanford University, 401 Quarry Rd, MC: 5797, Stanford, CA 94305 (amitetkin@stanford.edu).

Despite substantive evidence supporting the efficacy of antidepressants vs placebo for major depressive disorder (MDD),^{1,2} the lack of an accepted signature of placebo response has stymied the development of antidepressant drugs. As a result, trials in an otherwise heterogeneous sample of patients with MDD include many placebo responders, resulting in modest effect sizes (Cohen $d \sim 0.3$)^{1,3,4} and many failed trials. One avenue for improving the yield of drug development lies in grounding an understanding of the neural functioning associated with placebo response.

With 40% to 50% of patients with MDD not responding to antidepressant treatment,⁵⁻⁷ coupled with the low predictive value of clinical and sociodemographic variables,^{8,9} substantial attention has been directed at identifying pretreatment neural features that predict treatment response in MDD.^{6,7} Resting-state functional magnetic resonance imaging (fMRI) studies, for example, found that increased connectivity with the cognitive control network and decreased connectivity with the default mode network have been associated with antidepressant treatment response.^{6,7,10,11} However, prior work has not included a placebo control group, thus confounding treatment effects with nonspecific symptom change across time. Functional MRI is also ultimately limited by its cost and requirement for significant technical expertise.¹²

Electroencephalography (EEG), by contrast, provides a financially and logistically favorable neural assessment tool, with the additional benefits of its high temporal resolution and more direct measurement of neural function. With the use of EEG, increased frontal and parieto-occipital alpha power, increased anterior cingulate theta power, and greater global signal entropy have been found to be associated with better antidepressant treatment response.^{7,10,11,13-23} However, as with fMRI, the identification of EEG neural markers predictive of antidepressant response is limited by the lack of placebo-controlled clinical trials with sufficient statistical power.^{10,13} Moreover, one of the most promising treatment predictive findings from antidepressant treatment studies (namely, rostral cingulate theta power) appears to predict outcome for antidepressants and placebo.^{18,24} A recent meta-analysis suggests that, while there is great promise in EEG-guided MDD biomarker prediction, the field is currently limited by several factors, including sample size and lack of out-of-sample validation.¹⁹

Although EEG has great promise with respect to clinical translation, its low spatial precision and sensitivity to volume conduction (which leads to signal blurring) limit its utility for network-level connectivity analyses. This challenge has been addressed by recent advances in analytical methods, first in magnetoencephalography and then in EEG.^{25,26} By orthogonalizing source-estimated signals for instantaneous correlations, which arise primarily owing to volume conduction, and then correlating the power envelope (ie, instantaneous power) time series across different parts of the brain, studies have found large-scale connectivity patterns consistent with an fMRI-derived understanding of canonical human cortical networks.^{25,27}

We therefore sought to investigate how individual differences in EEG power envelope connectivity (PEC)-estimated

Key Points

Question What electroencephalographic connectivity features are neural moderators of antidepressant treatment response?

Findings In this secondary analysis of a randomized clinical trial, greater alpha-band and lower gamma-band connectivity—most prominently parietal—predicted better placebo treatment response and worse antidepressant treatment response. Lower connectivity levels in these moderating connections were associated with higher levels of anhedonia.

Meaning Key cortical features differentiating placebo response from antidepressant response were identified, providing an alternative direction toward establishing a placebo signature in clinical trials.

cortical networks in MDD differentially predicted outcome with an antidepressant vs placebo (ie, whether it could serve as a baseline moderator of treatment response). We did so using the largest neuroimaging-coupled, placebo-controlled randomized clinical trial in depression to date, to our knowledge—the Establishing Moderators and Biosignatures of Antidepressant Response in Clinic Care (EMBARC) study.²⁷ We calculated resting-state interregional PEC and graphed theoretical centrality measures to characterize EEG functional connectivity networks at baseline, and we examined within an intent-to-treat analytic framework how connectivity associated with treatment outcomes. Consistent with prior work,^{6,7,10,11} we hypothesized that greater PEC and node strength localized to the frontal and parietal cortices would predict better antidepressant treatment response specific to the antidepressant sertraline hydrochloride vs placebo.

Methods

Participants

This study is a nonprespecified secondary analysis of a randomized clinical trial. Additional details on study methods and participant characteristics are in eAppendix 1 and eTable 1 in the [Supplement](#) and in prior publications.^{18,27} Outpatients aged 18 to 65 years with a diagnosis of MDD²⁸ were recruited from July 29, 2011, to December 15, 2015, and assessed at 4 clinical sites (Columbia University, Massachusetts General Hospital, University of Michigan, and University of Texas Southwestern Medical Center). The study was approved by the institutional review boards of all sites, and participants provided written consent and received financial compensation. This study followed the Consolidated Standards of Reporting Trials (CONSORT) reporting guideline.

Assessments and Treatment Protocol

The EMBARC trial used a double-blind design, with randomization to 8 weeks of sertraline or placebo (N = 296; eFigure 1 in the [Supplement](#)). Randomization was stratified by site, depression severity, and chronicity using a block randomization procedure. The primary clinical outcome was total score on the 17-item Hamilton Rating Scale for Depression, administered at each study visit (baseline and weeks 1, 2, 3, 4, 6, and 8).

Electroencephalography

Each study site recorded data with a different EEG system and channel-space montage (eTable 2 in the [Supplement](#)). All data were sampled at 250 Hz, over 60 to 128 channels, depending on the site. Variables corresponding to site differences, including their multiplicative interaction with all other terms in the linear mixed models, were controlled for to reduce the possibility of site effects on results (see Statistical Analysis subsection; eTable 3 in the [Supplement](#)). Electroencephalography was performed during 8 total minutes of resting-state activity by alternating between 2-minute blocks of eyes-open (REO) and eyes-closed (REC) conditions.¹⁸

Analyses

A total of 268 treatment-randomized patients underwent EEG. Of these, 221 (82.5%) were included in the analyses (217 of whom had both REO and REC conditions), excluding 47 data sets because of poor EEG data quality. Of these patients, 106 were randomized to receive sertraline and 115 were randomized to receive placebo (eFigure 1 in the [Supplement](#)). Final data included 217 REC data sets (105 sertraline and 112 placebo) and 221 REO data sets (106 sertraline and 115 placebo) (see eTable 5 in the [Supplement](#) for participant identifiers included in the analyses). See eAppendix 1 in the [Supplement](#) for EEG preprocessing and analysis details. A repeated k-fold cross validation analysis was conducted and reported in eTable 12 in the [Supplement](#). Receiver operator characteristics (area under the curve), odds ratios, and effect sizes are reported for all analyses in eTable 13 in the [Supplement](#). Analyses controlling for effects of sex and age are reported in eTable 15 in the [Supplement](#).

Power Envelope Connectivity

All connectivity analyses were performed in source-space owing to the variability in montages across sites; therefore, each site's montage was retained. Source localization was performed in MATLAB 2014b (MathWorks Inc) using custom code for nonparametric minimum norm estimates.²⁹ All connectivity analyses were computed at the vertex level using 3003 vertices in MNI (Montreal Neurological Institute) template space, then averaged into 31 regions of interest (ROIs) in MNI space (eTable 4 in the [Supplement](#)). Power envelope connectivity²⁶ was derived from the following 4 canonical frequencies: theta (4.5-7.5 Hz), alpha (8-12 Hz), beta (12.5-30 Hz), and gamma (31-50 Hz). Further details can be found in eAppendix 1 and eFigure 2 in the [Supplement](#).

Node Strength Computation

The graph theoretical metric of node strength was computed for each ROI using MATLAB's implementation of the Brain Connectivity Toolbox.³⁰ Undirected node strength, quantified as the sum of weights of links connected to the node, was calculated on the orthogonalized PEC estimates.³¹

Statistical Analysis

Statistical analysis was performed from November 16, 2018, to May 23, 2019. Because our primary investigation was of the moderation of the linear trajectory of treatment outcome, we chose to perform prediction linear mixed-effects models, which

furthermore maintained an intent-to-treat framework. Linear mixed-effects models were applied to each ROI-to-ROI connectivity estimate and node strength measure.³²

Models predicted a linear trajectory of repeated 17-item Hamilton Rating Scale for Depression scores across time at baseline and week 1, 2, 3, 4, 6, and 8 (ie, end point) assessment points. Analyses were conducted using the NLME package in R, version 3.4.4 (R Foundation for Statistical Computing).³³ The brain metric \times treatment group \times time interaction, the primary term of interest, assessed differential symptom trajectories by treatment group as a function of EEG connectivity moderators (eTable 3 and eAppendix 1 in the [Supplement](#)). All *P* values were false discovery rate (FDR) corrected for multiple comparisons across all ROIs and frequency bands to control for type I errors (eAppendix 1 in the [Supplement](#)). All *P* values were from 2-sided tests and results were deemed statistically significant at an FDR-corrected $P \leq .0125$.

To further understand the clinical significance of treatment-moderating connectivity measures, we performed Pearson correlations of the FDR-significant connectivity and nodal strength effects with several clinical indices (correcting for multiple corrections across all correlations performed) across the entire sample. The clinical measures were the Childhood Trauma Questionnaire,³⁴ the Quick Inventory of Depressive Symptomatology,³⁵ the Mood and Anxiety Symptom Questionnaire,³⁶ the State-Trait Anxiety Questionnaire,³⁷ and the Snaith-Hamilton Pleasure Scale,³⁸ as well as the duration of the current depressive episode. *P* values extracted from the correlational analyses between clinical scales and connectivity features were FDR-corrected for multiple comparisons simultaneously across all pairwise comparisons of all scales and FDR-significant connectivity features. A supplementary analysis was conducted to assess connectivity features that change from baseline to 1 week into treatment (eTable 14, eFigure 3, and eAppendix 2 in the [Supplement](#)).

Results


A total of 221 participants underwent EEG recordings and had high-quality pretreatment EEG data. Participants included in the analyses were aged 18 to 65 years (150 women; mean [SD] age, 37.8 [12.7] years). The group mean (SD) score on the 17-item Hamilton Rating Scale for Depression was 18.35 (4.58) at baseline and 26.14 (30.37) across all time points.

PEC Moderators of Treatment Outcome

Initial Identification of Moderators

After correcting for multiple comparisons across all pairwise connectivity features and frequency bands, significant moderation effects for the REC condition were identified at 40 ROI-to-ROI pairwise power envelope connections using the alpha carrier frequency and at 31 connections using the gamma carrier frequency (eTable 6 in the [Supplement](#); **Figure 1B** and **C**). The regions associated with these connections were primarily frontal, parietal, temporal, and visual and somatosensory (**Figure 1B** and **C**).

Figure 1. Cortical Regions of Interest (ROIs), Alpha Power Envelope Connectivity (PEC), and Gamma PEC



A, A total of 31 ROIs (right and left ROIs are numbered the same; 14 bilateral ROIs [28 total] and 3 midline ROIs) were defined in Montreal Neurological Institute space, derived from independent components analysis parcellation of resting-state functional magnetic resonance imaging connectivity from 38 participants.²⁶
 B, Resting eyes-closed PEC between left mid-temporal gyrus and right angular gyrus moderation of outcome with sertraline (SER) vs placebo (PLA) treatment within the alpha band. C, Resting eyes-closed PEC between left anterior mid-frontal gyrus and right mid-temporal gyrus moderation of outcome with SER vs PLA treatment within the gamma band. Left panel: z scores for significantly moderating ROIs are represented in the ROI × ROI matrix plots. Middle panel: summed cortical connectivity (z scores) at each significantly moderating ROI. Right panel: visualization of moderation results reveals that significant prediction of treatment only in the PLA group was probably the cause of the moderation results. Shown are model-predicted 17-item Hamilton Rating Scale for Depression (HAMD17) values for the PLA and SER groups using an arbitrary median split on PEC for visualization purposes only. Low = below-median connectivity and high = above-median connectivity. The 2 ROIs comprising the pairwise connectivity feature visualized are outlined in red on the cortical images. AMFG indicates anterior mid-frontal gyrus; ANG, angular gyrus; DACC, dorsal anterior cingulate cortex; FEF, frontal eye fields; IFJ, inferior frontal junction; INS, insular cortex; IPL, inferior parietal lobe; IPS, inferior parietal sulcus; L, left; MPFC, medial prefrontal cortex; MTG, mid-temporal gyrus; ORB, orbitofrontal cortex; PCC, posterior cingulate cortex; PMFG, posterior mid-frontal gyrus; R, right; SEF, superior eye fields; SMC, sensorimotor cortex; SUP, supramarginal gyrus; and VI, bilateral primary visual cortex.

A total of 24 connections within the alpha, gamma, and beta frequencies significantly moderated treatment effect within the REO condition. Alpha-band moderators were primarily within the visual and parietal regions, whereas beta-band moderators were specific to parietal and frontal regions, and gamma-band moderators within the temporal, parietal, anterior cingulate, and visual and somatosensory cortices (eTable 7 in the Supplement; Figure 2).

Examining Moderation as a Function of Within-Group Treatment

To next understand whether these significant moderators were due to prediction of outcome in the sertraline group, placebo group, or both groups, we performed mixed models for each group separately. Because this analysis was performed to investigate what was causing the connectivity \times time \times treatment group interactions, we constrained our analyses only to FDR-significant connections (FDR-corrected $P \leq .0125$) identified as moderating in the full analysis. As such, we did not correct again for multiple comparisons because it was performed for visualization purposes only. We found that the moderating connections within the REC condition from the alpha and gamma carrier frequency analyses significantly predicted outcome primarily in the placebo group, with only a few predicting outcome in the sertraline group (eTable 6 in the Supplement). Within the REO condition, while the significantly predictive features across alpha, beta, and gamma carrier frequencies were predominately within the placebo group, most of these features were also predictive within the sertraline group (eTable 7 in the Supplement). Thus, REC connectivity moderators predominantly reflected placebo-driven effects, whereas those in the REO condition reflected effects from both groups.

Visualization of Within-Group Moderation Effects

To visualize treatment-moderating connections within the REC condition (with no further statistical analyses to avoid “double dipping”), we divided the sample with depression using a median split for each of the ROI-to-ROI PEC values found to significantly moderate treatment outcome (Figure 1B and C and Figure 2). This visualization revealed that greater alpha-band PEC within parietal, temporal, and visual regions predicted better treatment outcome with placebo (ie, steeper symptom change slopes) and worse treatment outcome with sertraline. In addition, reduced gamma-band PEC within the frontal, visual, somatomotor, parietal, and temporal regions also predicted better treatment outcomes with placebo and worse treatment outcomes with sertraline (Figure 1).²⁶

Within the REO condition, better outcomes in the placebo group and worse treatment outcomes in the sertraline group were predicted by greater alpha-band PEC within the visual and parietal regions; reduced gamma-band PEC within the frontal, temporal, parietal, anterior cingulate, and visual and somatosensory regions; and reduced beta-PEC within the frontal and parietal regions (Figure 2).

Power Envelope Node Strength Moderators of Treatment Outcome

As a complement to the connection-wise analyses, we next conducted the same linear mixed model analyses on node-level

connectivity strength measures, which provide a summary metric of the overall connectivity of each ROI. Of the 31 ROIs and 4 frequency bands entered in the linear mixed-effect model analyses, 3 regions within the alpha-band REC condition survived FDR correction in the moderation analysis (brain \times treatment group \times site interaction; eTable 8 in the Supplement; Figure 3). These regions were the right inferior parietal lobe, the right angular gyrus, and the right supramarginal gyrus. As in the analyses, visualizing the results using a median split, we found that a higher alpha-band nodal strength predicted better treatment outcome with placebo (Figure 3). Alpha-band nodal strength did not significantly predict treatment outcome with sertraline.

Association Between Clinical Severity Measures and PEC

We ran correlations between questionnaire data and FDR-significant connectivity pairs from the analyses. Correlations with the Snaith-Hamilton Pleasure Scale, which measures anhedonia,³⁹ survived correction. Baseline anhedonia ratings across the entire sample were negatively associated with alpha- and gamma-band connectivity metrics within the REC condition (eTable 9 in the Supplement; Figure 4A and B). The Snaith-Hamilton Pleasure Scale was also negatively associated with alpha node strength in the right inferior parietal lobe, the angular gyrus, and the supramarginal gyrus (Figure 4C; eTable 10 in the Supplement). In addition, greater symptom severity on the Quick Inventory of Depressive Symptomatology was positively associated with greater baseline gamma PEC within the frontal, parietal, temporal, insular, and cingulate cortical regions within the REC condition (eTable 11 in the Supplement). No REO connectivity measures survived correction.


Discussion

Here we establish pretreatment resting-state EEG connectivity features that moderated the treatment effect of the antidepressant sertraline vs placebo in a large randomized clinical trial in outpatients with MDD. Moderation effects were largely associated with outcomes in the placebo group and, to a lesser extent, in the sertraline group, a subset of which were associated with anhedonia and depressive symptom severity.

Neural Moderators of Treatment Response


The PEC measures found to significantly moderate treatment outcome were predominantly within the alpha-band, gamma-band, and, to a lesser extent, beta-band carrier frequencies. The synchronization of slow fluctuations in the power envelope of the band-limited oscillatory signal has been suggested to underlie large-scale functional network dynamics in both fMRI and magnetoencephalography.⁴⁰⁻⁴² Further supporting these findings, entraining the power envelopes of regions within a given resting-state network has been found to strengthen fMRI connectivity within the targeted networks.⁴³ Prior work has supported a negative association between alpha or beta oscillations and neural

Figure 2. Resting Eyes-Open Power Envelope Connectivity (PEC) Moderation of Outcome With Sertraline (SER) vs Placebo (PLA) Treatment Within the Alpha, Beta, and Gamma Bands



A, Alpha PEC between left and right primary visual cortex. B, Beta PEC between medial prefrontal cortex and left inferior parietal sulcus. C, Gamma PEC between left inferior parietal lobe and right primary visual cortex. Left panel: z scores for significantly moderating regions of interest (ROIs) are represented in the ROI × ROI matrix plots. Middle panel: summed cortical connectivity (z scores) at each significantly moderating ROI; ROIs with greater summed z scores are those with greater total moderating connectivity. Right panel: visualization of moderation results reveals that significant prediction of treatment only in the PLA group was probably the cause of the moderation results. Shown are model-predicted 17-item Hamilton Rating Scale for Depression (HAM-D17) values for the PLA and SER groups using an arbitrary median split on PEC for visualization purposes only. Low = below-median connectivity, high = above-median connectivity. Two ROIs comprising the pairwise connectivity feature visualized are outlined in red on the cortical images. AMFG indicates anterior mid-frontal gyrus; ANG, angular gyrus; DACC, dorsal anterior cingulate cortex; FEF, frontal eye fields; IFJ, inferior frontal junction; INS, insular cortex; IPL, inferior parietal sulcus; L, left; MPFC, medial prefrontal cortex; MTG, mid-temporal gyrus; ORB, orbitofrontal cortex; PCC, posterior cingulate cortex; PMFG, posterior mid-frontal gyrus; R, right; SEF, superior eye fields; SMC, sensorimotor cortex; SUP, supramarginal gyrus; and VI, bilateral primary visual cortex.

Figure 3. Alpha Power Envelope Connectivity (PEC) Node Strength Moderation of Outcome With Sertraline (SER) vs Placebo (PLA) Treatment



Left panel: z scores for significantly moderating features are represented in the above matrix plots. Right angular gyrus (RANG), right inferior parietal lobe (RIPL), and right supramarginal gyrus (RSUP) node strength significantly moderated treatment outcome within the alpha carrier frequency. Middle panel: summed cortical connectivity (z scores) at each significantly moderating region of interest. Right panel: visualization of RANG (outlined in red on cortical image) moderation results reveals that significant prediction of treatment only in the PLA group was probably the cause of the moderation results. Shown are model-predicted 17-item Hamilton Rating Scale for Depression (HAMDI7) values

for the PLA and SER groups using an arbitrary median split on PEC for visualization purposes only. Low = below-median connectivity, high = above-median connectivity. AMFG indicates anterior mid-frontal gyrus; DACC, dorsal anterior cingulate cortex; FEF, frontal eye fields; IFJ, inferior frontal junction; INS, insular cortex; IPS, inferior parietal sulcus; L, left; MPFC, medial prefrontal cortex; MTG, mid-temporal gyrus; ORB, orbitofrontal cortex; PCC, posterior cingulate cortex; PMFG, posterior mid-frontal gyrus; R, right; SEF, superior eye fields; SMC, sensorimotor cortex; and VI, bilateral primary visual cortex.

activation, as well as a positive association between gamma oscillations and neural activation, as measured by the fMRI blood oxygen level-dependent signal⁴⁴⁻⁴⁶ and intracranial recordings.⁴⁷⁻⁴⁹ See further discussion regarding band specificity in PEC in eAppendix 2 in the [Supplement](#).


The alpha-band connectivity features significantly moderating treatment response were predominantly between the parietal, visual, and temporal regions. Parietal alpha-band power has been heavily associated with reward anticipation⁵⁰⁻⁵² and has been shown to be reduced in patients with depression at rest and during reward anticipation.⁵³⁻⁵⁵ In addition, reduction in frontoparietal alpha power has been associated with reduced reward anticipation in depression.⁵⁵ Gamma-band connectivity was also found to significantly moderate treatment outcome, specific to the frontal, parietal, and sensorimotor regions. Some prior work has implicated gamma oscillations as a state-related marker of depression.^{56,57} Reductions in gamma-band power, complexity, and connectivity in the frontal-parietal cortices have been found in depression,⁵⁸ with gamma oscillation power increases with treatment correlating with improved treatment outcome.^{59,60} However, the relevance of band power to band-limited PEC is unknown and was not examined in these studies. Moreover, to our knowledge, no prior work examined the relevance of alpha-band or gamma-band power to placebo clinical outcome. As such, our findings provide important new insights in an area that has received relatively little research. The effect of these findings is further underscored by the size of the EMBARC trial relative to prior studies, which either lacked placebo groups or had samples 5 to 10 times smaller than those in the EMBARC trial.

Neural Predictors of Antidepressant Treatment Response

Placebo treatment can reduce depressive symptoms, so much so that it has become an increasing barrier to medication effectiveness research during the past few decades.^{1,4,61-69} This difficulty separating antidepressant from placebo treatment responses has motivated treatment prediction and moderation research, with the EMBARC trial being the largest such study using neuroimaging to date. Although some studies have found baseline neural features predictive of treatment response, most did not find these features to differentiate between antidepressant and placebo response.^{18,70-72} Moreover, because most used an open-label design without a placebo group, we still lack knowledge regarding the neural basis of response to placebo vs antidepressant medication.

Although all the significant features predicting treatment response were associated with placebo, a number of connectivity measures predicted antidepressant outcome in the opposite direction as placebo outcome. These measures were predominantly alpha-band connectivity within the parietal cortex (REC) and visual cortex (REO), beta-band connectivity within the frontal and parietal regions (REO), and gamma-band connectivity within and between the parietal, temporal, and visual and somatosensory cortices (REC and REO). This subset of features in particular may help identify individuals who will experience greater antidepressant and worse placebo responses, whereas the other measures would inform only placebo prediction. One known factor delineating placebo vs treatment responders is depressive symptom severity.⁷³ However, the connectivity features predictive of sertraline treatment response were not associated with the baseline metrics of Quick Inventory of Depressive Symp-

Figure 4. Association Between Electroencephalography Connectivity Moderators and Anhedonia



A, Alpha power envelope connectivity (PEC) between the left middle temporal gyrus and right inferior angular gyrus. B, Gamma PEC between left angular gyrus and left inferior parietal sulcus (see eTable 7 in the Supplement for statistics). C, Greater anhedonia (as measured by the Snaith-Hamilton Pleasure Scale [SHAPS]) is associated with lower alpha PEC node strength in the right angular gyrus (see eTable 8 in the Supplement for statistics). Two regions of interest (ROIs) comprising the pairwise connectivity feature, or single ROI comprising the node strength feature, are visualized are outlined in red on the cortical images. Diagonal lines indicate same ROI-to-ROI connections.

tomatology depressive symptom severity, reducing the possibility that these features simply reflect symptoms.

Neural Predictors of Placebo Treatment Response

Although a number of studies have examined the neurobiology of placebo response, only a handful have examined the neurobiological substrates of placebo response in depression.^{67,71,72} This work has also focused mostly on neural changes associated with placebo antidepressant re-

sponses, rather than pretreatment outcome prediction. The primary study within the literature investigating the baseline functional connectivity predictive of placebo treatment response identified increased rostral anterior cingulate cortex connectivity within the salience network as predictive of placebo treatment response.⁷²

The clinical significance of the treatment-moderating connectivity is further suggested by their association with

anhedonia, whereby greater neural connectivity and nodal strength was associated with reduced anhedonia. Anhedonia, or the loss of pleasure in activities, is a prominent symptom in depression.⁷⁴ Previous research has shown an association between anhedonia and disrupted neural reward processing.^{75,76} We speculate therefore that parietal PEC, in particular to the temporal and visual regions within the alpha band, may reflect in part the action of reward processing systems, which in turn predicts placebo response.

Limitations

This study has some limitations. The mechanism by which alpha-band and gamma-band features differentially influence treatment response requires further investigation. Future work will be needed to understand how pretreatment EEG connectivity predicts longer-term outcomes beyond 8 weeks as well as clinically important events such as relapse. The effect sizes were small for most of the significant features found to mod-

erate treatment outcome and, therefore, call for replication. Although both anhedonia and depressive severity were significantly associated with resting-state EEG treatment moderators, they explained only approximately 5% to 10% of variance in connectivity.

Conclusions

Our findings provide insight into the neural connectivity features that are specifically predict placebo response in depression, differentially from sertraline treatment response, potentially involving a role for anhedonia. From a treatment perspective, capitalizing on the therapeutic components leading to placebo response differentially from antidepressant response could provide an alternative direction toward clinical treatment of patients with depression.⁶⁸

ARTICLE INFORMATION

Accepted for Publication: October 5, 2019.

Published Online: January 2, 2020.

doi:10.1001/jamapsychiatry.2019.3867

Author Affiliations: Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, California (Rolle, Fonzo, Wu, Toll, Etkin); Wu Tsai Neuroscience Institute, Stanford University, Stanford, California (Rolle, Fonzo, Wu, Toll, Etkin); Veterans Affairs Palo Alto Healthcare System, Palo Alto, California (Rolle, Fonzo, Wu, Toll, Etkin); Sierra Pacific Mental Illness, Research, Education, and Clinical Center, Palo Alto, California (Rolle, Fonzo, Etkin); Department of Psychiatry, Dell Medical School, The University of Texas at Austin (Fonzo); School of Automation Science and Engineering, South China University of Technology, Guangzhou, Guangdong, China (Wu); Department of Psychiatry, University of Texas Southwestern Medical Center, Dallas (Jha, Cooper, Chin-Fatt, Trombello, Trivedi); Department of Psychiatry, Harvard Medical School, Boston, Massachusetts (Pizzagalli, Deckersbach, Fava); New York State Psychiatric Institute, Department of Psychiatry, College of Physicians and Surgeons of Columbia University, New York (Weissman); now at Alto Neuroscience Inc, Los Altos, California (Etkin).

Author Contributions: Ms Rolle had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Drs Trivedi and Etkin contributed equally to this research.

Concept and design: Pizzagalli, Trivedi, Etkin.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Rolle, Deckersbach, Etkin.

Critical revision of the manuscript for important intellectual content: Rolle, Fonzo, Wu, Toll, Jha, Cooper, Chin-Fatt, Pizzagalli, Trombello, Fava, Weissmann, Trivedi, Etkin.

Statistical analysis: Rolle, Fonzo, Wu, Toll, Chin-Fatt, Trivedi, Etkin.

Obtained funding: Weissmann, Trivedi.

Administrative, technical, or material support:

Fonzo, Toll, Jha, Cooper, Pizzagalli, Trombello, Fava, Weissmann, Etkin.

Supervision: Toll, Cooper, Pizzagalli, Trombello, Trivedi, Etkin.

Conflict of Interest Disclosures: Dr Pizzagalli reported receiving funding during the last 3 years from the National Institute of Mental Health, the Dana Foundation, and Brain and Behavior Research Foundation; and receiving consulting fees or honoraria from Akili Interactive Labs, Alkermes, BlackThorn Therapeutics, Boehringer Ingelheim, Compass, and Posit Science for activities unrelated to the current research. Dr Trombello reported currently owning stock in Merck and Gilead Sciences, both of which are unrelated to the current project. Dr Deckersbach reported receiving research funding from the National Institutes of Health, the National Institute of Mental Health, National Alliance for Research on Schizophrenia and Depression, Tourette Syndrome Association, International OCD (Obsessive Compulsive Disorder) Foundation, Tufts University, Depressive and Bipolar Disorder Alternative Treatment Foundation, and Otsuka Pharmaceuticals; receiving honoraria, consultation fees, and/or royalties from the Massachusetts General Hospital Psychiatry Academy, BrainCells Inc, Clintara LLC Inc, Systems Research and Applications Corp, Boston University, the Catalan Agency for Health Technology Assessment and Research, the National Association of Social Workers Massachusetts, the Massachusetts Medical Society, Tufts University, the National Institute on Drug Abuse, the National Institute of Mental Health, Oxford University Press, Guilford Press, and Rutledge; participating in research funded by Defense Advanced Research Projects Agency, the National Institutes of Health, National Institute on Aging, the Agency for Healthcare Research and Quality, the Patient-Centered Outcomes Research Institute, Janssen Pharmaceuticals, The Forest Research Institute, Shire Development Inc, Medtronic, Cyberonics, Northstar, and Takeda. Dr Fava reported receiving research support from Abbott Laboratories, Alkermes Inc, American Cyanamid, Aspect Medical Systems, AstraZeneca, Avanir Pharmaceuticals, BioResearch, BrainCells Inc, Bristol-Myers Squibb, CeNeRx BioPharma, Cephalon, Clintara LLC, Cerecor, Covance, Covidien, Eli Lilly and Company, EnVivo Pharmaceuticals Inc,

Euthymics Bioscience Inc, Forest Pharmaceuticals Inc, Ganeden Biotech Inc, GlaxoSmithKline, Harvard Clinical Research Institute, Hoffman-LaRoche, Icon Clinical Research, i3 Innovus/Ingenix, Janssen R&D LLC, Jed Foundation, Johnson & Johnson Pharmaceutical Research & Development, Lichtwer Pharma GmbH, Lorex Pharmaceuticals, Lundbeck Inc, MedAvante, Methylation Sciences Inc, National Alliance for Research on Schizophrenia & Depression, National Center for Complementary and Alternative Medicine, National Institute of Drug Abuse (NIDA), National Institute of Mental Health, Neuralstem Inc, Novartis AG, Organon Pharmaceuticals, PamLab LLC, Pfizer Inc, Pharmacia-Upjohn, Pharmaceutical Research Associates Inc, Pharmavite LLC, Pharmorx Therapeutics, Photothera, Reckitt Benckiser, Roche Pharmaceuticals, RCT Logic LLC (formerly Clinical Trials Solutions LLC), Sanofi-Aventis US LLC, Shire, Solvay Pharmaceuticals Inc, Stanley Medical Research Institute, Synthelabo, Tal Medical, and Wyeth-Ayerst Laboratories; serving as advisor or consultant to Abbott Laboratories, Acadia, Affectis Pharmaceuticals AG, Alkermes Inc, Amarin Pharma Inc, Aspect Medical Systems, AstraZeneca, Auspex Pharmaceuticals, Avanir Pharmaceuticals, AXSOME Therapeutics, Bayer AG, Best Practice Project Management Inc, Biogen, BioMarin Pharmaceuticals Inc, Biovail Corporation, BrainCells Inc, Bristol-Myers Squibb, CeNeRx BioPharma, Cephalon Inc, Cerecor, CNS Response Inc, Compellis Pharmaceuticals, Cypress Pharmaceutical Inc, DiagnoSearch Life Sciences (P) Ltd, Dinippon Sumitomo Pharma Co Inc, Dov Pharmaceuticals Inc, Edgemont Pharmaceuticals Inc, Eisai Inc, Eli Lilly and Company, EnVivo Pharmaceuticals Inc, ePharmaSolutions, EPIX Pharmaceuticals Inc, Euthymics Bioscience Inc, Fabre-Kramer Pharmaceuticals Inc, Forest Pharmaceuticals Inc, Forum Pharmaceuticals, GenOmind LLC, GlaxoSmithKline, Grunenthal GmbH, i3 Innovus/Ingenix, Intracellular, Janssen Pharmaceuticals, Jazz Pharmaceuticals Inc, Johnson & Johnson Pharmaceutical Research & Development LLC, Knoll Pharmaceuticals Corp, Labopharm Inc, Lorex Pharmaceuticals, Lundbeck Inc, MedAvante Inc, Merck & Co Inc, MSI Methylation Sciences Inc, Naurex Inc, Nestle Health Sciences, Neuralstem Inc, Neuronetics Inc, NextWave Pharmaceuticals,

Novartis AG, Nutrition 21, Orexigen Therapeutics Inc, Organon Pharmaceuticals, Osmotica, Otsuka Pharmaceuticals, PamLab LLC, Pfizer Inc, PharmaStar, Pharmavite LLC, Pharmorx Therapeutics, Precision Human Biotechnology, Prexa Pharmaceuticals Inc, Puretech Ventures, PsychoGenics, Psylin Neurosciences Inc, RCT Logic LLC (formerly Clinical Trials Solutions LLC), Rexahn Pharmaceuticals Inc, Ridge Diagnostics Inc, Roche, Sanofi-Aventis US LLC, Sepracor Inc, Servier Laboratories, Schering-Plough Corporation, Solvay Pharmaceuticals Inc, Somaxon Pharmaceuticals Inc, Somerset Pharmaceuticals Inc, Sunovion Pharmaceuticals, Supernus Pharmaceuticals Inc, Synthelabo, Taisho Pharmaceutical, Takeda Pharmaceutical Company Limited, Tal Medical Inc, Tetrigenex Pharmaceuticals Inc, TransForm Pharmaceuticals Inc, Transcept Pharmaceuticals Inc, Vanda Pharmaceuticals Inc, and VistaGen; receiving speaking or publishing fees from Adamed Co, Advanced Meeting Partners, American Psychiatric Association, American Society of Clinical Psychopharmacology, AstraZeneca, Belvoir Media Group, Boehringer Ingelheim GmbH, Bristol-Myers Squibb, Cephalon Inc, CME Institute/Physicians Postgraduate Press Inc, Eli Lilly and Company, Forest Pharmaceuticals Inc, GlaxoSmithKline, Immedex LLC, MGH Psychiatry Academy/Primedia, MGH Psychiatry Academy/Reed Elsevier, Novartis AG, Organon Pharmaceuticals, Pfizer Inc, PharmaStar, United BioSource Corp and Wyeth-Ayerst Laboratories; having equity holdings in Compellis and PsyBrain Inc; holding a patent for Sequential Parallel Comparison Design, which is licensed by MGH to Pharmaceutical Product Development LLC; having a patent application for a combination of ketamine plus scopolamine in major depressive disorder, licensed by MGH to Biohaven; receiving copyright royalties for the MGH Cognitive & Physical Functioning Questionnaire, Sexual Functioning Inventory, Antidepressant Treatment Response Questionnaire, Discontinuation-Emergent Signs & Symptoms, Symptoms of Depression Questionnaire, and SAFER; and receiving publishing royalties of books from Lippincott Williams & Wilkins, Wolters Kluwer, and World Scientific Publishing Co Pte Ltd. Dr Weissman reported receiving funding in the past 2 years from the National Institute of Mental Health, the National Institute on Drug Abuse, the National Alliance for Research on Schizophrenia and Depression, the Sackler Foundation, and the Templeton Foundation; and receiving royalties from Oxford University Press, Perseus Press, the American Psychiatric Association Press, and MultiHealth Systems. Dr Trivedi reported serving as an advisor or consultant and receiving fees from (lifetime disclosure) Abbott Laboratories Inc, Abdi Ibrahim, Akzo (Organon Pharmaceuticals Inc), Alkermes, AstraZeneca, Axon Advisors, Bristol-Myers Squibb Company, Cephalon Inc, Cerecor, CME Institute of Physicians, Concert Pharmaceuticals Inc, Eli Lilly & Company, Evotec, Fabre Kramer Pharmaceuticals Inc, Forest Pharmaceuticals, GlaxoSmithKline, Janssen Global Services LLC, Janssen Pharmaceutica Products LP, Johnson & Johnson PRD, Libby, Lundbeck, Meade Johnson, MedAvante, Medtronic, Merck, Mitsubishi Tanabe Pharma Development America Inc, Naurex, Neuronetics, Otsuka Pharmaceuticals, PamLab, Parke-Davis Pharmaceuticals Inc, Pfizer Inc, PgxHealth, Phoenix Marketing Solutions, Rexahn Pharmaceuticals, Ridge Diagnostics, Roche

Products Ltd, Sepracor, SHIRE Development, Sierra, SK Life and Science, Sunovion, Takeda, Tal Medical/Puretech Venture, Targacept, Transcept, VantagePoint, Vivus, and Wyeth-Ayerst Laboratories; and receiving grants and research support from the Agency for Healthcare Research and Quality, Cyberonics Inc, the National Alliance for Research in Schizophrenia and Depression, the National Institute of Mental Health, and the National Institute on Drug Abuse. Dr Etkin (lifetime disclosure) reported receiving salary and equity from Alto Neuroscience Inc; holding equity in Mindstrong Health, Akili Interactive, and Sizing for unrelated work; receiving research funding from the National Institute of Mental Health, Department of Veterans Affairs, Cohen Veterans Bioscience, Brain and Behavior Research Foundation, Dana Foundation, Brain Resource Inc, and the Stanford Neurosciences Institute; and serving as a consultant for Cervel, Takeda, Posit Science, Acadia, Otsuka, Lundbeck, and Janssen. No other disclosures were reported.

Funding/Support: The EMBARC study was supported by the National Institute of Mental Health of the National Institutes of Health under awards U01MH092221 (Dr Trivedi) and U01MH092250 (Dr Weissman). This work was also funded in part by the Hersh Foundation (Dr Trivedi, principal investigator). Drs Wu and Etkin were funded by National Institutes of Health grant DP1 MH116506. Dr Wu was funded by the National Key Research and Development Plan of China (grant 2017YFB1002505) and the National Natural Science Foundation of China (grants 61876063 and 61836003). Ms Rolle was funded by the National Science Foundation, Graduate Research Fellowship Program (grant 2016180976).

Role of the Funder/Sponsor: The National Institutes of Health had a role in the design and conduct of the study, but had no role in collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; nor decision to submit the manuscript for publication.

REFERENCES

- Cipriani A, Furukawa TA, Salanti G, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Lancet*. 2018; 391(10128):1357-1366. doi:10.1016/S0140-6736(17)32802-7
- Khan A, Faucett J, Lichtenberg P, Kirsch I, Brown WA. A systematic review of comparative efficacy of treatments and controls for depression. *PLoS One*. 2012;7(7):e41778. doi:10.1371/journal.pone.0041778
- Fournier JC, DeRubeis RJ, Hollon SD, et al. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA*. 2010; 303(1):47-53. doi:10.1001/jama.2009.1943
- Kirsch I, Deacon BJ, Huedo-Medina TB, Scoboria A, Moore TJ, Johnson BT. Initial severity and antidepressant benefits: a meta-analysis of data submitted to the Food and Drug Administration. *PLoS Med*. 2008;5(2):e45. doi:10.1371/journal.pmed.0050045
- Trivedi MH, Rush AJ, Wisniewski SR, et al; STAR*D Study Team. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry*. 2006;163(1):28-40. doi:10.1176/appi.ajp.163.1.28
- Dichter GS, Gibbs D, Smoski MJ. A systematic review of relations between resting-state functional-MRI and treatment response in major depressive disorder. *J Affect Disord*. 2015;172:8-17. doi:10.1016/j.jad.2014.09.028
- Harmer CJ. Neural predictors of treatment response in depression. *Curr Behav Neurosci Rep*. 2014;1(3):125-133. doi:10.1007/s40473-014-0021-2
- Bagby RM, Ryder AG, Crispien C. Psychosocial and clinical predictors of response to pharmacotherapy for depression. *J Psychiatry Neurosci*. 2002;27(4):250-257.
- Nierenberg AA. Predictors of response to antidepressants: general principles and clinical implications. *Psychiatr Clin North Am*. 2003;26(2):345-352. doi:10.1016/S0193-953X(02)00105-3
- Kemp AH, Gordon E, Rush AJ, Williams LM. Improving the prediction of treatment response in depression: integration of clinical, cognitive, psychophysiological, neuroimaging, and genetic measures. *CNS Spectr*. 2008;13(12):1066-1086. doi:10.1017/S1092852900017120
- Pizzagalli DA. Frontocingulate dysfunction in depression: toward biomarkers of treatment response. *Neuropsychopharmacology*. 2011;36(1):183-206. doi:10.1038/npp.2010.166
- Carmichael O, Schwarz AJ, Chatham CH, et al. The role of fMRI in drug development. *Drug Discov Today*. 2018;23(2):333-348. doi:10.1016/j.drudis.2017.11.012
- Olbrich S, Arns M. EEG biomarkers in major depressive disorder: discriminative power and prediction of treatment response. *Int Rev Psychiatry*. 2013;25(5):604-618. doi:10.3109/09540262.2013.816269
- Arns M, Bruder G, Hegerl U, et al. EEG alpha asymmetry as a gender-specific predictor of outcome to acute treatment with different antidepressant medications in the randomized iSPOT-D study. *Clin Neurophysiol*. 2016;127(1):509-519. doi:10.1016/j.clinph.2015.05.032
- Frodl T. Recent advances in predicting responses to antidepressant treatment. *F1000Res*. 2017;6:6. doi:10.12688/f1000research.10300.1
- Jaworska N, Wang H, Smith DM, Blier P, Knott V, Protzner AB. Pre-treatment EEG signal variability is associated with treatment success in depression. *Neuroimage Clin*. 2017;17:368-377. doi:10.1016/j.nicl.2017.10.035
- Pizzagalli D, Pascual-Marqui RD, Nitschke JB, et al. Anterior cingulate activity as a predictor of degree of treatment response in major depression: evidence from brain electrical tomography analysis. *Am J Psychiatry*. 2001;158(3):405-415. doi:10.1176/appi.ajp.158.3.405
- Pizzagalli DA, Webb CA, Dillon DG, et al. Pretreatment rostral anterior cingulate cortex theta activity in relation to symptom improvement in depression: a randomized clinical trial. *JAMA Psychiatry*. 2018;75(6):547-554. doi:10.1001/jamapsychiatry.2018.0252
- Widge AS, Bilge MT, Montana R, et al. Electroencephalographic biomarkers for treatment response prediction in major depressive illness: a meta-analysis. *Am J Psychiatry*. 2019;176(1):44-56. doi:10.1176/appi.ajp.2018.17121358

20. Wade EC, Iosifescu DV. Using electroencephalography for treatment guidance in major depressive disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2016;1(5):411-422. doi:10.1016/j.bpsc.2016.06.002
21. Olbrich S, van Dinteren R, Arns M. Personalized medicine: review and perspectives of promising baseline EEG biomarkers in major depressive disorder and attention deficit hyperactivity disorder. *Neuropsychobiology*. 2015;72(3-4):229-240. doi:10.1159/000437435
22. Schiller MJ. Quantitative electroencephalography in guiding treatment of major depression. *Front Psychiatry*. 2019;9:779. doi:10.3389/fpsy.2018.00779
23. Mumtaz W, Malik AS, Yasin MAM, Xia L. Review on EEG and ERP predictive biomarkers for major depressive disorder. *Biomed Signal Process Control*. 2015;22:85-98. doi:10.1016/j.bspc.2015.07.003
24. Korb AS, Hunter AM, Cook IA, Leuchter AF. Rostral anterior cingulate cortex theta current density and response to antidepressants and placebo in major depression. *Clin Neurophysiol*. 2009;120(7):1313-1319. doi:10.1016/j.clinph.2009.05.008
25. Hipp JF, Hawellek DJ, Corbetta M, Siegel M, Engel AK. Large-scale cortical correlation structure of spontaneous oscillatory activity. *Nat Neurosci*. 2012;15(6):884-890. doi:10.1038/nn.3101
26. Toll R, Wu W, Naparstek S, et al. An electroencephalography connectomic profile of post-traumatic stress disorder. *Am J Psychiatry*. In press.
27. Trivedi MH, McGrath PJ, Fava M, et al. Establishing Moderators and Biosignatures of Antidepressant Response in Clinical Care (EMBARC): rationale and design. *J Psychiatr Res*. 2016;78:11-23. doi:10.1016/j.jpsychires.2016.03.001
28. First M, Spitzer R, Gibbon M, William J. *Structured Clinical Interview for DSM-IV-TR Axis I disorders, Research Version, Patient Edition (SCID-I/P)*. New York: New York State Psychiatric Institute, Biometric Research; 2002.
29. Hämäläinen MS, Ilmoniemi RJ. Interpreting magnetic fields of the brain: minimum norm estimates. *Med Biol Eng Comput*. 1994;32(1):35-42. doi:10.1007/BF02512476
30. Rubinov M, Sporns O. Complex network measures of brain connectivity: uses and interpretations. *Neuroimage*. 2010;52(3):1059-1069. doi:10.1016/j.neuroimage.2009.10.003
31. Fallani FdeV, Costa LdaF, Rodriguez FA, et al. A graph-theoretical approach in brain functional networks: possible implications in EEG studies. *Nonlinear Biomed Phys*. 2010;4(suppl 1):S8. doi:10.1186/1753-4631-4-S1-S8
32. Lindstrom ML, Bates DM. Nonlinear mixed effects models for repeated measures data. *Biometrics*. 1990;46(3):673-687. doi:10.2307/2532087
33. Pinheiro J, Bates D, DebRoy S, Sarkar D; R Core Team. nlme: Linear and nonlinear mixed effects models. R package version 3.1-142 [computer program]. <https://cran.r-project.org/web/packages/nlme/index.html>. Accessed November 22, 2019.
34. Bernstein DP, Fink L. *Childhood Trauma Questionnaire: A Retrospective Self-Report Manual*. San Diego, CA: Psychological Corporation; 1998.
35. Rush AJ, Trivedi MH, Ibrahim HM, et al. The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol Psychiatry*. 2003;54(5):573-583. doi:10.1016/S0006-3223(02)01866-8
36. Watson D, Clark LA. *The Mood and Anxiety Symptom Questionnaire*. Iowa City: University of Iowa; 1991.
37. Spielberger CD, Gorsuch RL, Lushene RE. *Manual for the State-Trait Anxiety Inventory (Self-Evaluation Questionnaire)*. Palo Alto, CA: Consulting Psychologists Press; 1970.
38. Snaith RP, Hamilton M, Morley S, Humayan A, Hargreaves D, Trigwell P. A scale for the assessment of hedonic tone the Snaith-Hamilton Pleasure Scale. *Br J Psychiatry*. 1995;167(1):99-103. doi:10.1192/bjp.167.1.99
39. Nakonezny PA, Morris DW, Greer TL, et al. Evaluation of anhedonia with the Snaith-Hamilton Pleasure Scale (SHAPS) in adult outpatients with major depressive disorder. *J Psychiatr Res*. 2015;65:124-130. doi:10.1016/j.jpsychires.2015.03.010
40. de Pasquale F, Della Penna S, Snyder AZ, et al. Temporal dynamics of spontaneous MEG activity in brain networks. *Proc Natl Acad Sci U S A*. 2010;107(13):6040-6045. doi:10.1073/pnas.0913863107
41. Liu Z, Fukunaga M, de Zwart JA, Duyn JH. Large-scale spontaneous fluctuations and correlations in brain electrical activity observed with magnetoencephalography. *Neuroimage*. 2010;51(1):102-111. doi:10.1016/j.neuroimage.2010.01.092
42. Nikouline VV, Linkenkaer-Hansen K, Huttunen J, Ilmoniemi RJ. Interhemispheric phase synchrony and amplitude correlation of spontaneous beta oscillations in human subjects: a magnetoencephalographic study. *Neuroreport*. 2001;12(11):2487-2491. doi:10.1097/00001756-200108080-00040
43. Bächinger M, Zerbi V, Moisa M, et al. Concurrent tACS-fMRI reveals causal influence of power synchronized neural activity on resting state fMRI connectivity. *J Neurosci*. 2017;37(18):4766-4777. doi:10.1523/JNEUROSCI.1756-16.2017
44. Hall EL, Robson SE, Morris PG, Brookes MJ. The relationship between MEG and fMRI. *Neuroimage*. 2014;102(pt 1):80-91. doi:10.1016/j.neuroimage.2013.11.005
45. Mukamel R, Gelbard H, Arieli A, Hasson U, Fried I, Malach R. Coupling between neuronal firing, field potentials, and fMRI in human auditory cortex. *Science*. 2005;309(5736):951-954. doi:10.1126/science.1110913
46. Zumer JM, Brookes MJ, Stevenson CM, Francis ST, Morris PG. Relating BOLD fMRI and neural oscillations through convolution and optimal linear weighting. *Neuroimage*. 2010;49(2):1479-1489. doi:10.1016/j.neuroimage.2009.09.020
47. Haegens S, Nacher V, Luna R, Romo R, Jensen O. α -Oscillations in the monkey sensorimotor network influence discrimination performance by rhythmical inhibition of neuronal spiking. *Proc Natl Acad Sci U S A*. 2011;108(48):19377-19382. doi:10.1073/pnas.1117910108
48. Hayden BY, Smith DV, Platt ML. Electrophysiological correlates of default-mode processing in macaque posterior cingulate cortex. *Proc Natl Acad Sci U S A*. 2009;106(14):5948-5953. doi:10.1073/pnas.0812035106
49. Ossandón T, Jerbi K, Vidal JR, et al. Transient suppression of broadband gamma power in the default-mode network is correlated with task complexity and subject performance. *J Neurosci*. 2011;31(41):14521-14530. doi:10.1523/JNEUROSCI.2483-11.2011
50. Gorka SM, Phan KL, Shankman SA. Convergence of EEG and fMRI measures of reward anticipation. *Biol Psychol*. 2015;112:12-19. doi:10.1016/j.biopsycho.2015.09.007
51. Liu F, Guo W, Yu D, et al. Classification of different therapeutic responses of major depressive disorder with multivariate pattern analysis method based on structural MR scans. *PLoS One*. 2012;7(7):e40968. doi:10.1371/journal.pone.0040968
52. Wisniewski D, Reverberi C, Momennejad I, Kahnt T, Haynes JD. The role of the parietal cortex in the representation of task-reward associations. *J Neurosci*. 2015;35(36):12355-12365. doi:10.1523/JNEUROSCI.4882-14.2015
53. Nelson BD, Perlman G, Klein DN, Kotov R, Hajcak G. Blunted neural response to rewards as a prospective predictor of the development of depression in adolescent girls. *Am J Psychiatry*. 2016;173(12):1223-1230. doi:10.1176/appi.ajp.2016.15121524
54. Shankman SA, Klein DN, Tenke CE, Bruder GE. Reward sensitivity in depression: a biobehavioral study. *J Abnorm Psychol*. 2007;116(1):95-104. doi:10.1037/0021-843X.116.1.95
55. Shankman SA, Nelson BD, Sarapas C, et al. A psychophysiological investigation of threat and reward sensitivity in individuals with panic disorder and/or major depressive disorder. *J Abnorm Psychol*. 2013;122(2):322-338. doi:10.1037/a0030747
56. Fitzgerald PJ, Watson BO. Gamma oscillations as a biomarker for major depression: an emerging topic. *Transl Psychiatry*. 2018;8(1):177. doi:10.1038/s41398-018-0239-y
57. Fitzgerald PB, Laird AR, Maller J, Daskalakis ZJ. A meta-analytic study of changes in brain activation in depression. *Hum Brain Mapp*. 2008;29(6):683-695. doi:10.1002/hbm.20426
58. Akar SA, Kara S, Agambayev S, Bilgic V. Nonlinear analysis of EEG in major depression with fractal dimensions. *Conf Proc IEEE Eng Med Biol Soc*. 2015;2015:7410-7413. doi:10.1109/EMBC.2015.7320104
59. Noda Y, Zomorodi R, Saeki T, et al. Resting-state EEG gamma power and theta-gamma coupling enhancement following high-frequency left dorsolateral prefrontal rTMS in patients with depression. *Clin Neurophysiol*. 2017;128(3):424-432. doi:10.1016/j.clinph.2016.12.023
60. Pathak Y, Salami O, Baillet S, Li Z, Butson CR. Longitudinal changes in depressive circuitry in response to neuromodulation therapy. *Front Neural Circuits*. 2016;10:50. doi:10.3389/fncir.2016.00050
61. Enck P, Klosterhalfen S, Weimer K, Horing B, Zipfel S. The placebo response in clinical trials: more questions than answers. *Philos Trans R Soc Lond B Biol Sci*. 2011;366(1572):1889-1895. doi:10.1098/rstb.2010.0384
62. Stolk P, Ten Berg MJ, Hemels ME, Einarson TR. Meta-analysis of placebo rates in major depressive disorder trials. *Ann Pharmacother*. 2003;37(12):1891-1899. doi:10.1345/aph.1D172
63. Walsh BT, Seidman SN, Sysko R, Gould M. Placebo response in studies of major depression:

- variable, substantial, and growing. *JAMA*. 2002;287(14):1840-1847. doi:10.1001/jama.287.14.1840
64. Al-Harbi KS. Treatment-resistant depression: therapeutic trends, challenges, and future directions. *Patient Prefer Adherence*. 2012;6:369-388. doi:10.2147/PPA.S29716
65. Cipriani A, Geddes JR. Placebo for depression: we need to improve the quality of scientific information but also reject too simplistic approaches or ideological nihilism. *BMC Med*. 2014;12:105. doi:10.1186/1741-7015-12-105
66. Khan A, Redding N, Brown WA. The persistence of the placebo response in antidepressant clinical trials. *J Psychiatr Res*. 2008;42(10):791-796. doi:10.1016/j.jpsychores.2007.10.004
67. Peciña M, Bohnert AS, Sikora M, et al. Association between placebo-activated neural systems and antidepressant responses: neurochemistry of placebo effects in major depression. *JAMA Psychiatry*. 2015;72(11):1087-1094. doi:10.1001/jamapsychiatry.2015.1335
68. Rutherford BR, Roose SP. A model of placebo response in antidepressant clinical trials. *Am J Psychiatry*. 2013;170(7):723-733. doi:10.1176/appi.ajp.2012.12040474
69. Salanti G, Chaimani A, Furukawa TA, et al. Impact of placebo arms on outcomes in antidepressant trials: systematic review and meta-regression analysis. *Int J Epidemiol*. 2018;47(5):1454-1464. doi:10.1093/ije/dyy076
70. Leuchter AF, Cook IA, Witte EA, Morgan M, Abrams M. Changes in brain function of depressed subjects during treatment with placebo. *Am J Psychiatry*. 2002;159(1):122-129. doi:10.1176/appi.159.1.122
71. Mayberg HS, Silva JA, Brannan SK, et al. The functional neuroanatomy of the placebo effect. *Am J Psychiatry*. 2002;159(5):728-737. doi:10.1176/appi.ajp.159.5.728
72. Sikora M, Heffernan J, Avery ET, Mickey BJ, Zubieta JK, Peciña M. Salience network functional connectivity predicts placebo effects in major depression. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2016;1(1):68-76. doi:10.1016/j.bpsc.2015.10.002
73. Khan A, Brodhead AE, Kolts RL, Brown WA. Severity of depressive symptoms and response to antidepressants and placebo in antidepressant trials. *J Psychiatr Res*. 2005;39(2):145-150. doi:10.1016/j.jpsychores.2004.06.005
74. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Washington, DC: American Psychiatric Association; 2013.
75. Der-Avakian A, Markou A. The neurobiology of anhedonia and other reward-related deficits. *Trends Neurosci*. 2012;35(1):68-77. doi:10.1016/j.tins.2011.11.005
76. Heshmati M, Russo SJ. Anhedonia and the brain reward circuitry in depression. *Curr Behav Neurosci Rep*. 2015;2(3):146-153. doi:10.1007/s40473-015-0044-3